



# 22<sup>nd</sup> INTERNATIONAL DIATOM SYMPOSIUM

Ghent, Belgium 26-31 August 2012

## ABSTRACTS

22nd International Diatom Symposium

Ghent, Belgium

26-31 August 2012

VLIZ SPECIAL PUBLICATION 58

Organized by:

Lab. Protistology & Aquatic Ecology  
Biology Department  
Ghent University  
Krijgslaan 281, S8  
B-9000 Ghent, Belgium  
[www.pae.ugent.be](http://www.pae.ugent.be)

National Botanic Garden of Belgium  
Domein of Bouchout  
Nieuwelaan 38  
B-1860 Meise, Belgium  
[www.Botanicgarden.be](http://www.Botanicgarden.be)

## **ORGANIZING COMMITTEE**

Koen Sabbe, Bart Van de Vijver, Wim Vyverman, Annick Verween, Koenraad Muylaert, Annick Wilmotte, Jean-Pierre Descy, Kateřina Kopalová, Rebecca van der Wiele, Pieter Vanormelingen, Caroline Souffreau

## **SCIENTIFIC COMMITTEE**

Koen Sabbe (Ghent University, Belgium)  
Wim Vyverman (Ghent University, Belgium)  
Bart Van de Vijver (National Botanic Garden of Belgium)  
Koenraad Muylaert (KU Leuven, KULAK, Belgium)  
Marie Huysman (Ghent University, Belgium)  
Peter Kroth (University of Konstanz, Germany)  
Eileen Cox (Natural History Museum London, UK)  
Regine Jahn (Botanischer Garten und Botanisches Museum Berlin-Dahlem, Germany)  
Angela Wulff (Goteborgs Universitet, Sweden)  
Frédéric Rimet (French National Institute for Agricultural Research INRA, France)  
Shigeki Mayama (Tokyo Gakugei University, Japan)  
Sarah A. Spaulding (University of Colorado, Boulder, USA)

### **This publication should be cited as follows:**

Sabbe Koen, Van de Vijver Bart & Vyverman Wim (Eds). 2012 Twentysecond International Diatom Symposium, Aula Academica, Ghent, 26-31 August 2012. Abstracts. VLIZ Special Publication 58, 260p.

Reproduction is authorized, provided that appropriate mention is made of the source.

ISSN 1377-0950

## THE ACTION OF TERAHERTZ LASER IRRADIATION ON DIATOM FRUSTULES

Vadim V. Annenkov<sup>1</sup>, Tatjana N. Basharina<sup>1</sup>, Alexander S. Kozlov<sup>2</sup>, Elena N. Danilovtseva<sup>1</sup> & Alexander K. Petrov<sup>2</sup>

<sup>1</sup>Limnological Institute of the Siberian Branch of the Russian Academy of Sciences

<sup>2</sup>Institute of Chemical Kinetics and Combustion

The main feature of diatoms is the presence of a silica exoskeleton and a lot of biochemical and molecular biology studies are focused on the elucidation of the biosilica synthesis mechanism. A diatom cell uses a special organelle SDV (Silica Deposition Vesicle) to obtain elements of new diatom frustules. This organelle is relatively large, visible in light microscopy but there are no reliable data concerning the structure of its membrane and its contents. The silica exoskeleton is the main obstacle in a soft destruction of diatom cells and the isolation of non-destroyed SDVs. Silica frustules are highly resistant to mechanical and chemical actions, so searching for new approaches in order to remove silica frustules from the diatom cells is an actual task.

Laser ablation using a free electron laser (FEL) is a novel method for soft destruction of supramolecular systems stabilized by hydrogen bonds [A. K. Petrov et al. Nucl. Instrum. Methods Phys. Res., Sect. A. 575:68]. The advantage of ablation with FEL consists in correspondence between laser frequencies ( $40\text{-}80\text{ cm}^{-1}$ ) and vibrations of hydrogen bonds, and so the destruction of covalent bonds is unlikely. The softness of the ablation technique has been confirmed by the retention of enzymatic activity of horseradish peroxidase and PCR-proven integrity of lambda phage DNA, plasmid pUC18DNA after all manipulations.

We have found that the action of terahertz laser irradiation on the fresh-water diatom *Synedra acus* subsp. *radians* (Kützing) Skabichevskii results in the splitting of diatom frustules. The most effective was an irradiation of a wave length of  $100\text{ }\mu\text{m}$  which is close to the diatom length. It seems as silica valves are torn away from the cell membrane and the whole cell contents remains in a non-destroyed inner membrane. Chloroplasts and SDVs do not change their shape after the valve removing.

The study was partially supported by the Siberian Branch of the Russian Academy of Sciences (project #39).